

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12P 13/00, 7/64, C07F 9/10	A1	(11) International Publication Number: WO 00/52190 (43) International Publication Date: 8 September 2000 (08.09.00)
(21) International Application Number: PCT/US00/04589 (22) International Filing Date: 23 February 2000 (23.02.00) (30) Priority Data: 09/259,603 1 March 1999 (01.03.99) US (71) Applicant: KEMIN INDUSTRIES, INC. [US/US]; 2100 Maury Street, Des Moines, IA 50301-0070 (US). (72) Inventors: SAS, Benedikt; Steenweg Op Ravels 209, B-2360 Oud-Turnhout (BE). PEYS, Eric; Ossenberg 81, B-2490 Balen (BE). (74) Agent: KENT, A., Herink; Davis, Brown, Koehn, Shors & Roberts, P.C., The Financial Center, Suite 2500, 666 Walnut Street, Des Moines, IA 50309-3993 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: IMPROVED METHOD FOR THE CONVERSION OF LECITHIN INTO LYSOLECITHIN (57) Abstract Conversion of crude lecithin is performed by incubating the lecithin with a lipase-phospholipase mixture in water/polyol environment. The incubation is carried out at about 40-50 °C and a pH of about 7 to about 8 while stirring continuously. The reaction simultaneously produces lysolipid, lysophospholipid, monoglycerides, and diglycerides. The conversion rate of lecithin is more than 80 % based on the ration of LPC/PC (lysophosphatidylcholine/phosphatidylcholine). Calcium chloride can be added to the reaction mixture to capture released fatty acids.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

IMPROVED METHOD FOR THE CONVERSION OF LECITHIN INTO LYSOLECITHIN

Background of the Invention

The present invention relates generally to lecithin conversion. More particularly, this invention relates to an improved method of enzymatic conversion of lecithin to produce lysolecithin. Even more particularly, this invention relates to a method of simultaneously producing lysolipid, lysophospholipid, and mono- and diglycerides from lecithin.

Lecithin and lysolecithin have been used in various industries as emulsifiers. These emulsifiers are used in a variety of products, e.g., foods, cosmetics, drugs, nutritional supplements, and chemicals.

Enzymatic conversion of phosphatidylcholine to lysophosphatidylcholine has been known for years. Previous researchers have determined that several factors such as pH, temperature, and surface concentration of the molecules affect this conversion. Additionally, presence or absence of particular ions, such as calcium, or solvents have been determined to affect the conversion process.

These prior art processes have low conversion rates and are unable to produce a cost-effective process.

The process of the current invention converts lecithin at a higher conversion rate while using lecithin, rather than phosphatidylcholine, and producing a desired, product containing lysolipids, lysophospholipids, and mono- and diglycerides.

Summary of the Invention

Accordingly, it is an object of the present invention provide an improved method for converting lecithin into lysolecithin.

Another object of the invention is to provide a method which simultaneously converts lipids and phospholipids of crude lecithin into lysolipids, lysophospholipids, and mono- and diglycerides.

Another object of the invention is to provide a method which simultaneously converts lipids and phospholipids of crude lecithin into lysolipids, lysophospholipids, and mono- and diglycerides at a greater conversion percentage than prior art methods.

The invention is a method for the improved conversion of lipids and phospholipids present in crude lecithin into their lyso- forms and the simultaneous production of mono- and diglycerides to get an improved emulsifier. This conversion is performed by means of an enzymatic treatment. The enzymatic treatment consists of the hydrolysis and/or alcoholysis of lipids and phospholipids in a water/polyol mixture, by means of a blend of lipase and phospholipase A2. The lipase cleaves ester bonds of the lipids and produces glycerol and mono- and diglycerides. The phospholipase A2 cleaves the ester bond of the phospholipids at the C2 position of the glycerol backbone. The resultant lysophospholipids are surface-active amphiphiles that contain only one long chain fatty acyl group and a large polar head group consisting of a free hydroxyl on the glycerol backbone and an anionic or zwitterionic moiety bonded to the same backbone via a phosphate ester. Common moieties include choline, ethanolamine, and inositol.

The conversion is performed by incubating lecithin with the lipase-phospholipase mixture in an aqueous/polyol environment with or without the addition of calcium chloride (CaCl_2) to capture the released fatty acids. The lecithin is mixed

with water and polyols at 40-50 °C that has been made slightly basic. This mixture is kept at 50°C and is continuously stirred. The pH of this mixture is continuously monitored and kept at pH 7 to 8 by addition of a base. When the lecithin and solvent are well-mixed the lipase-phospholipase is added. The conversion is checked by means of an HPLC method of analysis.

Until now the conversion of phospholipids of crude lecithin into lysophospholipids with phospholipase A2 on a large scale resulted in a low conversion percentage. The present method is able to increase the conversion percentage while simultaneously producing mono- and diglycerides. The conversion is more than 80% based on the ratio of LPC/(LPC+PC) (LPC=lysophosphatidylcholine; PC=phosphatidylcholine).

Brief Description of the Drawings

Figure 1 is a high-performance liquid chromatography- evaporative light-scattering detector (HPLC-ELSD) chromatogram of crude lecithin.

Figure 2 is a HPLC-ELSD chromatogram of the product of Example 1.

Figure 3 is a HPLC-ELSD chromatogram of the product of Example 2.

Figure 4 is a HPLC-ELSD chromatogram of the product of Example 3.

Detailed Description of the Invention

The invention is an improved method for simultaneously producing lysolipids, lysophospholipids, and mono- and diglycerides from crude lecithin. The method produces a high conversion rate of over 80% based on the ratio of LPC/PC (lysophosphatidylcholine/phosphatidylcholine) in a single step.

Crude lecithin can be used in the process of the current invention. Prior art processes generally use phosphatidylcholine. Lecithin is available from various commercial sources.

The lecithin is added to a water/polyol mixture. The water/polyol is in a 6 to 1 weight ratio. The lecithin is added to the mixture in a 3 to 7 weight ratio. In a preferred embodiment of the invention, the water/polyol mixture is at about 40°C to about 50 °C and has been made slightly basic. The preferred polyol for the process is glycerol. Though this is the preferred polyol, any polyol may be used. The pH of the mixture is preferred to be about 7 to 8. The range of pH that will work depends on the workable pH range of both lipase and phospholipase A2. The pH may be adjusted using a base. A base that has been used is sodium hydroxide (NaOH) at 10N. Any base may be used as long as it does not interfere with the desired reactions. It is desired to keep the combination of lecithin/water/polyol at a constant pH. A pH-stat has been used to achieve a constant pH. The combination of lecithin/water/polyol is preferably continuously stirred.

A blend of lipase and phospholipase A2 is added to the lecithin/water/polyol combination. The lipase and phospholipase A2 enzymes are available from various commercial sources. A lipase/phospholipase A2 mixture from Lovesgrove Research Ltd. has been used. The conversion can be performed with any other combination of lipase and phospholipase A2 as long as there is an overlap in their workable pH range. The activities of the enzymes used in the examples was about 16 units per gram of crude lecithin for the lipase and about 48 units per gram of crude lecithin for the phospholipase A2. One lipase unit is equal to the amount of enzyme that liberates one micromole of fatty acids per minute from tricaprylin under the conditions of the assay

(pH=7.5, 40°C). One phospholipase unit is equal to the amount of enzyme that liberates one micromole of fatty acids per minute from soy phosphatidylcholine under the conditions of the assay (pH=7.5, 40°C). Though the full range of amounts and ratios of these enzymes that will be effective have not been tested, half the amount of enzymes was effective, albeit over a longer incubation period. This is incubated for a length of time necessary to achieve a desired conversion rate. This length of time is dependent upon the amount and activity of the enzymes added. The end point can be found by checking with HPLC, or by watching the pH-stat data for leveling. The pH-stat data of the preferred embodiment showed conversion completed after approximately 12 hours.

Certain enzymes may require calcium (Ca^{2+}) ions to exert their activity. Calcium chloride (CaCl_2) can be added to the mixture, though it is not necessary. Calcium chloride is available from various commercial sources. Calcium ions catch the fatty acids that are released during the conversion. Any compound which serves this function but does not interfere with the desired results may be used. The calcium chloride can be added at an amount in moles from zero up to half the number of moles of fatty acids expected to be released during incubation.

The methods of this invention are further illustrated by the following experimental examples.

EXAMPLES

Example 1

Four hundred twenty grams (420 g) of water and 70 g glycerol were mixed. The pH of the mixture was adjusted to 7.5 with NaOH 10N. This mixture was stirred

at approximately 100 rpm and kept at a temperature of 50°C. The pH was kept constant at 7.5 with NaOH 10N. A lipase-phospholipase A2 enzyme mixture was added. Two hundred ten grams (210 g) crude lecithin was added, and the mixture was incubated for 18 hours.

Results: Table 1 shows the results of Examples 1-3. There was 3.3% LPC of (LPC+PC) in the crude lecithin. The resulting product of this procedure contained 93.4% LPC of (LPC+PC). See Figures 1 and 2 for the chromatograms of the starting and ending materials.

Figure 1. HPLC-ELSD Chromatogram of starting material (crude lecithin).

Figure 2. HPLC-ELSD Chromatogram of the final product using the above procedure of Example 1.

Example 2

Three thousand grams (3000 g) water and 500 g glycerol were mixed. The pH of the mixture was adjusted to 7.5 using NaOH 10N. This mixture was stirred at approximately 100 rpm and kept at 50°C. The pH was constantly kept at 7.5 using NaOH 10N. A lipase-phospholipase A2 mixture was added. Fifteen hundred grams (1500 g) crude lecithin was added, and the mixture was incubated for 21 hours.

Results: Table 1 shows the results of Examples 1-3. The crude lecithin starting material was the same as that of Example 1, i.e., containing 3.3% LPC of (LPC+PC). The resulting product of this procedure contained 95.6% LPC of (LPC+PC). See Figures 1 and 3 for the chromatograms of the starting and ending materials.

Figure 3. HPLC-ELSD Chromatogram of the final product using the above procedure of Example 2.

Example 3

Two thousand two hundred twenty grams (2220 g) water and 370 g glycerol were mixed, and the pH of the mixture was adjusted to 7.5 with NaOH 10N. This mixture was stirred at approximately 100 rpm and kept at 50°C. The pH was kept constant using NaOH 10N. A lipase-phospholipase A2 enzyme mixture was added. One thousand one hundred ten grams (1110 g) crude lecithin was added, and the mixture was incubated for 21 hours.

Results: Table 1 shows the results of Examples 1-3. The crude lecithin starting material was the same as that of Example 1, i.e., containing 3.3% LPC of (LPC+PC). The resulting product of this procedure contained 94.4% LPC of (LPC+PC). See Figures 1 and 4 for the chromatograms of the starting and ending materials. Figure 4. HPLC-ELSD Chromatogram of the final product using the above procedure of Example 3.

TABLE 1

Material	%LPC of (LPC+PC)
Crude Lecithin	3.3
Lovesgrove	18
Example 1 Product	93.4
Example 2 Product	95.6
Example 3 Product	94.4

Example 4

A competitor's modified lecithin was compared against lecithin converted by the process of the current invention using both P-NMR and the HPLC-ELSD methods. The results are given in Table 2. The Lovesgrove modified lecithin is made via a bacterial/enzymatic conversion using phospholipase A2 (see UK and PCT patent No. 9205014.5 and U.S. Patent 5,759,537).

TABLE 2

Wt. %	Crude Lecithin	Lovesgrove Research Limited modified lecithin (³¹ P-NMR data)	Lovesgrove Research Limited modified lecithin (HPLC data)	Converted Lecithin of Examples 1-3 (HPLC data)
PA (phosphatidic acid)	3.21	4.82	3.01	
PE (phosphatidylethanolamine)	7.51	10.08	8.19	
PI (phosphatidylinositol)	6.87	7.93	6.88	1.4 ± 0.4
LPA+LPE (lysophosphatidic acid and lysophosphatidylethanolamine)		1.77	1.90	5.3 ± 0.7
PC (phosphatidylcholine)	12.24	12.32	11.09	0.43 ± 0.10
LPC (lysophosphatidylcholine)	0.42	1.99 (1- LPC + 2- LPC)	1.19	7.27 ± 0.23
%LPC of (LPC+PC)	3.3	13.91	9.69	93.4-95.6%

Although the invention has been described with respect to a preferred embodiment thereof, it is to be also understood that it is not to be so limited since

changes and modifications can be made therein which are within the full intended scope of this invention as defined by the appended claims.

What is claimed is:

1. A method for simultaneously producing lysolipids, lysophospholipids, monoglycerides, and diglycerides comprising:
 - a. combining lecithin with a water/polyol mixture to form a combination,
 - b. contacting the combination with a blend of lipase and phospholipase A2,
 - c. incubating for a length of time effective to achieve the desired conversion rate.
2. The method of claim 1 further comprising adding calcium chloride to the lecithin/water/polyol combination.
3. The method of claim 2 wherein the calcium chloride is added in a range of about zero moles to about half the number of moles of fatty acids expected to be released during incubation.
4. The method of claim 1 wherein the weight ratio of water to polyol is about 6 to about 1.
5. The method of claim 1 wherein the water/polyol mixture pH is slightly basic.
6. The method of claim 5 wherein the pH is about 7 to about 8.
7. The method of claim 6 wherein the pH is about 7.5.


8. The method of claim 1 wherein the water/polyol mixture is maintained at a temperature of about 40°C to about 50°C.
9. The method of claim 1 wherein the weight ratio of lecithin to water/polyol is about 3 to about 7.
10. The method of claim 1 wherein the combination of lecithin/water/polyol is continuously stirred.
11. The method of claim 1 wherein the combination of lecithin/water/polyol is kept at a pH of about 7 to about 8.
12. The method of claim 11 wherein the pH is about 7.5.
13. The method of claim 1 wherein the polyol is glycerol.
14. A method for simultaneously producing lysolipid and lysophospholipid and mono- and diglycerides comprising:
 - a. combining lecithin with a slightly basic water/polyol mixture at a temperature of about 40°C to about 50°C to form a combination,
 - b. maintaining the combination at about 40°C to about 50°C,
 - c. stirring the combination continuously,
 - d. maintaining the pH of the combination at about 7 to about 8 using a base,

- e. adding a lipase-phospholipase A2 blend to the well-mixed combination,
 - f. incubating for a length of time effective to achieve the desired conversion rate.
15. The method of claim 14 wherein the temperature is about 50°C.
16. The method of claim 14 wherein the pH is about 7.5.
17. The method of claim 14 wherein the base is sodium hydroxide.
18. The method of claim 14 further comprising adding calcium chloride to the lecithin/water/polyol combination.
19. A method for simultaneously producing lysolipid and lysophospholipid and mono- and diglycerides comprising:
- a. combining lecithin with a water/glycerol to form a combination,
 - b. adjusting the pH of the combination to about 7.5 using sodium hydroxide,
 - c. stirring the combination continuously at approximately 100 rpm and at a temperature of about 50°C,
 - d. maintaining the combination at about 50°C,
 - e. maintaining the pH of the combination at about 7.5 using a sodium hydroxide,
 - f. adding a lipase-phospholipase A2 blend to the well-mixed combination,

g. incubating for approximately 12 hours.

20. The method of 19 further comprising adding calcium chloride to the lecithin/water/glycerol combination.

FIGURE 1






FIGURE 3





FIGURE 4



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/04589

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12P 13/00, 7/64; C07F 9/10

US CL : 435/128, 134, 198

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/128, 134, 198

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US 5,955,327 A (HIRAI et al) 21 September 1999, see entire disclosure.	1-20

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

11 MAY 2000

Date of mailing of the international search report

13 JUL 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DR. HERBERT J. LILLING

Telephone No. (703) 308-0196